Journal of Advances in Developmental Research (IJAIDR)

E-ISSN: 0976-4844 ● Website: www.ijaidr.com ● Email: editor@ijaidr.com

Optimization of Agronomic Practices for Enhancing Lemon Grass (Cymbopogon citratus) Yield in Saline Environments

Dr Dharmendra Singh

Department of Soil Science and Agriculture Chemistry, Dayanand College, Ajmer (Rajasthan)

Abstract

This review paper explores the optimization of agronomic practices for enhancing lemongrass (*Cymbopogon citratus*) yield in saline environments, particularly within the Indian context. Saline soils, prevalent in regions such as Rajasthan, Gujarat, and Tamil Nadu, present significant challenges to agricultural productivity. Effective management of soil fertility, irrigation, and water usage is crucial for overcoming the adverse effects of salinity. The paper discusses various strategies, including the application of organic and inorganic soil amendments (e.g., gypsum, farmyard manure), appropriate irrigation techniques (such as drip irrigation and leaching), and efficient water management practices. These techniques have been shown to improve soil health, reduce salt accumulation, and enhance plant growth, leading to better lemongrass yields. The review also highlights the role of nutrient management and crop-specific cultivation practices in mitigating salinity stress. The integration of modern technologies, such as soil moisture sensors and optimized fertilizer application, is essential for improving water use efficiency and overall crop productivity. This paper aims to provide a comprehensive understanding of agronomic practices that can help increase the sustainability and profitability of lemongrass cultivation in saline-prone areas of India, ensuring higher biomass and essential oil yields.

Keywords: Lemongrass, Saline soils, Agronomic practices, Water management, Drip irrigation, Soil amendments, Fertility management, Essential oil yield, Salinity stress, Indian agriculture

1. Introduction

Lemongrass (*Cymbopogon citratus*) is a tropical plant of considerable economic importance, known for its aromatic leaves used in the production of essential oils, which are widely utilized in the food, cosmetic, and pharmaceutical industries. Its agronomic characteristics, including its resilience to various environmental conditions, make it a viable crop in many regions, particularly in India. The plant is valued for its ability to thrive in diverse soil types, with notable benefits in producing high-quality oil for industrial applications (Chauhan et al., 2006). However, the cultivation of lemongrass in saline environments poses significant challenges due to the adverse effects of salt on plant growth, productivity, and essential oil yield.

Salinity is a major abiotic stress affecting crop production worldwide, particularly in arid and semi-arid regions of India. Approximately 6.7 million hectares of land in India are affected by salinity, with states like Rajasthan, Gujarat, and Tamil Nadu facing significant salinity issues in agricultural production (Yadav et al., 2007). In saline soils, the high concentration of soluble salts interferes with water

absorption by plant roots, leading to physiological stress. For lemongrass, this stress results in stunted growth, reduced leaf biomass, and ultimately lower essential oil content. According to Bhattacharyya et al. (2006), salinity concentrations exceeding 6 dS/m can drastically reduce lemongrass yield, with studies reporting a yield loss of up to 40% under such conditions.

Despite these challenges, research has demonstrated that lemongrass can be cultivated successfully in saline environments through the optimization of agronomic practices. Various studies suggest that the application of proper irrigation techniques, soil amendments, and nutrient management strategies can mitigate the detrimental effects of salinity. For instance, saline irrigation practices, when coupled with leaching and the addition of organic matter, have shown promising results in improving yield in saline conditions (Rattan et al., 2008). Moreover, the adoption of salt-tolerant varieties and improved management of water resources has led to yield improvements of up to 30% in certain saline-affected regions of India (Chauhan & Singh, 2009).

In conclusion, while the cultivation of lemongrass in saline environments presents several challenges, optimized agronomic practices offer a promising solution to enhance productivity. This review aims to explore these practices and provide a comprehensive analysis of their impact on lemongrass yield under saline conditions in the Indian context.

2. Effect of Salinity on Lemon Grass Growth

Salinity is a significant abiotic stress factor that negatively impacts the growth and productivity of many crops, including lemongrass (*Cymbopogon citratus*). In India, where around 6.7 million hectares of land are affected by salinity, particularly in states like Rajasthan and Gujarat, this issue is especially critical for the successful cultivation of various crops, including aromatic plants (Yadav et al., 2007). Lemongrass is sensitive to soil salinity, with higher salt concentrations leading to physiological disruptions such as reduced water uptake, osmotic stress, and nutrient imbalances (Rattan et al., 2008).

Research has shown that salinity levels exceeding 4 dS/m can cause a decline in the growth parameters of lemongrass, including leaf area, biomass production, and essential oil content (Chauhan et al., 2006). According to a study by Bhattacharyya et al. (2006), at a salinity level of 6 dS/m, the fresh weight of lemongrass plants was reduced by 30-40%, significantly lowering the overall yield. Salinity stress induces stomatal closure, limiting transpiration and reducing the plant's ability to carry out photosynthesis effectively, leading to poor growth and a decrease in essential oil yield (Munns, 2002).

Furthermore, the impact of salinity on lemongrass varies with its growth stages. During germination and early seedling development, high salinity levels have been shown to inhibit root development and germination rates. Bhattacharyya et al. (2006) observed that salinity levels of 8 dS/m completely hindered seed germination in lemongrass, making it challenging to establish crops in saline environments. During the vegetative and reproductive stages, salt stress leads to the accumulation of toxic ions such as sodium (Na^+) and chloride (Cl^-) in plant tissues, further impairing metabolic processes and growth (Chauhan & Singh, 2009).

Despite these adverse effects, lemongrass has shown some resilience to moderate salinity, especially in areas where saline irrigation practices are integrated with soil amendments. However, the degree of tolerance remains highly dependent on the specific salinity levels and the management practices adopted

(Rattan et al., 2008). Thus, understanding the specific thresholds of salinity tolerance in lemongrass is critical for developing strategies to optimize its cultivation in saline soils.

3. Agronomic Practices for Lemon Grass Cultivation

Successful cultivation of lemongrass (*Cymbopogon citratus*) in saline environments requires the optimization of several agronomic practices, aimed at mitigating the effects of soil salinity and enhancing crop yield. In India, where saline soil is prevalent in regions like Rajasthan, Gujarat, and Tamil Nadu, these practices become critical to ensuring sustainable and profitable lemongrass farming (Yadav et al., 2007). Effective management of soil and irrigation is essential to improving the plant's tolerance to salt stress and promoting higher productivity.

Soil Management and Amendments: The physical properties of saline soils, such as high electrical conductivity (EC), affect plant growth by disrupting water uptake and nutrient absorption. To counteract this, soil amendments such as organic matter, gypsum, and biochar are widely used to improve soil structure, enhance water infiltration, and reduce salt toxicity (Chauhan et al., 2006). For instance, the application of gypsum at a rate of 2-3 tons per hectare has been shown to improve soil salinity conditions and increase the yield of lemongrass by approximately 25-30% (Singh et al., 2008). Additionally, organic matter improves the cation-exchange capacity (CEC) of the soil, helping to bind harmful salts and reduce their availability to plants.

Irrigation Practices: The use of efficient irrigation methods, particularly drip irrigation, has proven beneficial in managing saline water and minimizing salt buildup around the root zone. Drip irrigation systems not only conserve water but also provide consistent moisture to the roots, reducing salt concentration in the soil (Rattan et al., 2008). Research indicates that the adoption of drip irrigation, combined with the leaching of excess salts, can increase the growth rate of lemongrass by up to 40% in saline environments (Chauhan & Singh, 2009). It is also recommended to implement periodic irrigation cycles to flush out accumulated salts, a practice that can significantly improve plant health.

Nutrient Management: Fertilizer management plays a crucial role in optimizing the growth of lemongrass in saline soils. Salinity often causes nutrient imbalances, particularly in nitrogen and potassium, which are essential for plant growth. Studies suggest that balanced fertilization with a mix of macro and micro-nutrients can offset these imbalances. Application of potassium nitrate (KNO3) has been shown to enhance salt tolerance in lemongrass, improving its biomass production and essential oil yield by 20-25% (Bhattacharyya et al., 2006).

In conclusion, the adoption of appropriate soil management techniques, efficient irrigation systems, and balanced nutrient practices is essential for optimizing lemongrass cultivation in saline environments. These agronomic practices not only improve plant growth and yield but also contribute to sustainable agricultural practices in regions with saline soils.

4. Impact of Fertilization on Lemon Grass Yield in Saline Conditions

Fertilization plays a pivotal role in enhancing the yield of lemongrass (*Cymbopogon citratus*) in saline environments. The availability and balance of nutrients are often disrupted in saline soils, which affects plant growth and essential oil production. The application of appropriate fertilizers, particularly those

that improve nutrient uptake and counteract salinity-induced nutrient imbalances, can significantly boost lemongrass productivity.

In saline conditions, the plant faces challenges in nutrient absorption due to high salt concentrations that hinder the movement of water and nutrients through the root system. Potassium, nitrogen, and phosphorus are particularly essential for lemongrass growth, as they directly influence photosynthesis, growth, and oil production. Studies suggest that the application of potassium-rich fertilizers, such as potassium nitrate (KNO3), can help mitigate the negative effects of salinity by enhancing the osmotic regulation of the plant and improving its ability to manage water stress (Bhattacharyya et al., 2006). Potassium also helps in reducing the sodium-to-potassium ratio in plant tissues, which is crucial for maintaining cell function under salt stress (Rattan et al., 2008).

Table 1: Impact of Fertilization on Lemongrass Yield under Saline Conditions

Fertilizer Type	Application Rate (kg/ha)	Fresh Biomass Yield (kg/ha)	Essential Oil Yield (L/ha)
Control (no fertilizer)	-	2,500	30
Nitrogen (N)	80	3,000	35
Potassium (K)	60	3,200	38
NPK (15-15-15)	100	3,500	40

Source: Singh et al., 2008

As shown in Table 1, the application of potassium significantly enhances both biomass and essential oil yield. Nitrogen also plays an important role in promoting plant growth, but potassium appears to be more effective in improving yield under saline stress. Additionally, the combined application of nitrogen, phosphorus, and potassium (NPK) fertilizers has shown the highest yield, both in terms of biomass and essential oil production.

Moreover, the timing and method of fertilizer application are crucial for optimizing yield. Fertilizer application during the vegetative phase, followed by periodic top-dressing, has been found to enhance the growth of lemongrass under salt stress. Balanced nutrient management, incorporating both macro and micro-nutrients, helps in alleviating the physiological stress caused by high salinity and ensures better nutrient availability to the plants (Chauhan & Singh, 2009).

In conclusion, the effective use of fertilizers, particularly potassium and NPK blends, can substantially increase lemongrass yield in saline soils. Fertilization strategies must be tailored to the specific needs of the crop and local soil conditions to maximize both growth and oil production.

5. Role of Irrigation in Optimizing Lemon Grass Yield in Saline Environments

Irrigation is a crucial factor for optimizing the growth and yield of lemongrass (*Cymbopogon citratus*) in saline environments. In regions with saline soils, the availability of fresh water is often limited, which poses significant challenges to crop production. Therefore, efficient irrigation techniques are necessary

to manage water resources and mitigate the adverse effects of salinity. In India, where regions like Rajasthan and Gujarat face high salinity, the implementation of proper irrigation practices becomes essential to ensure sustainable lemongrass farming.

Drip Irrigation: Drip irrigation is particularly effective in saline conditions as it delivers water directly to the root zone, reducing the chances of salt accumulation around the plant roots. This method helps in maintaining optimal moisture levels while minimizing the risk of salinity buildup in the soil (Chauhan et al., 2006). Drip irrigation has been shown to improve the yield of lemongrass by 20-40%, particularly in arid and semi-arid regions of India where water quality is often poor and saline (Singh et al., 2008).

Leaching Requirement: To combat soil salinity, leaching—flushing excess salts from the root zone—is an important aspect of irrigation management. Studies have indicated that the leaching requirement for lemongrass grown under saline conditions can range from 0.15 to 0.25, depending on the salinity level of the water used for irrigation (Yadav et al., 2007). The leaching process, when coupled with efficient irrigation methods, helps to maintain the soil's osmotic balance and reduces the toxicity of salts.

Table 2: Impact of Different Irrigation Methods on Lemongrass Yield under Saline Conditions

Irrigation Method	Water Application (L/plant/day)	Fresh Biomass Yield (kg/ha)	Essential Oil Yield (L/ha)
Surface Irrigation	25	2,800	32
Drip Irrigation	20	3,200	38
Furrow Irrigation	22	3,000	35

Source: Singh et al., 2008

As shown in Table 2, drip irrigation results in higher biomass and essential oil yield compared to traditional surface and furrow irrigation methods. This is primarily due to the efficient use of water and the prevention of salt buildup, which can hinder plant growth.

Water Quality and Salinity Management: In saline environments, the quality of water used for irrigation also plays a significant role. The Electrical Conductivity (EC) of irrigation water should ideally be below 2 dS/m for optimal lemongrass growth. Higher EC values result in reduced water uptake and plant stress, negatively impacting yield (Chauhan & Singh, 2009). Therefore, regular monitoring of water salinity is essential to maintain sustainable irrigation practices.

In conclusion, efficient irrigation techniques like drip irrigation, coupled with proper water management practices, can significantly enhance the growth and yield of lemongrass in saline environments. By minimizing salt accumulation and optimizing water use, farmers can achieve better productivity and sustainability.

6. Integrated Pest Management (IPM) Strategies for Lemon Grass in Saline Environments

Integrated Pest Management (IPM) is a sustainable approach that combines biological, cultural, mechanical, and chemical control methods to manage pests in crops. In saline environments, where environmental stressors impact plant health, implementing effective IPM strategies for lemongrass

(*Cymbopogon citratus*) is crucial to reduce the impact of pest infestations while maintaining environmental sustainability. In India, pest management in saline soils requires specialized strategies due to the unique challenges posed by salinity stress.

Cultural Practices: Cultural practices such as crop rotation, planting resistant varieties, and adjusting planting dates are integral components of IPM. By rotating lemongrass with other non-susceptible crops, pest populations can be disrupted, reducing the chances of pest buildup (Singh et al., 2007). Additionally, planting lemongrass in well-drained, raised beds can improve soil aeration, reducing the proliferation of soil-borne pests that thrive in saline soils.

Biological Control: The use of natural predators or biocontrol agents to manage pests is gaining traction in saline environments. For instance, the introduction of beneficial insects such as ladybird beetles and predatory mites can significantly reduce aphid populations, a common pest in lemongrass fields (Yadav et al., 2008). Additionally, the use of biopesticides such as neem oil, which is effective against a wide range of pests, can be an eco-friendly alternative to chemical pesticides.

Chemical Control: While chemical pesticides are sometimes necessary, they should be used sparingly to prevent resistance buildup and to protect the beneficial organisms in the ecosystem. In saline environments, the application of systemic insecticides with low toxicity to non-target species is recommended. These can be applied during the early stages of pest infestations when pest populations are still manageable (Chauhan & Sharma, 2009).

Table 3: Effectiveness of IPM Practices in Managing Pests in Lemongrass under Saline Conditions

IPM Strategy	Pest Reduction (%)	Biomass Yield (kg/ha)	Essential Oil Yield (L/ha)
Cultural Practices	25	3,000	34
Biological Control	30	3,200	37
Chemical Control	45	3,400	40
Integrated IPM (Cultural + Biological + Chemical)	65	3,800	45

Source: Yadav et al., 2008

As shown in Table 3, a combination of cultural, biological, and chemical methods under an integrated IPM approach has the highest impact on pest reduction, leading to significant improvements in both biomass and essential oil yield. The combined strategies reduce pest pressure effectively while maintaining crop health, especially in challenging saline environments.

In conclusion, implementing a holistic IPM strategy is essential for sustainable pest management in lemongrass cultivation under saline conditions. By integrating cultural practices, biological control, and judicious use of chemical agents, farmers can maintain high yields while minimizing the environmental impact of pesticide use.

7. Soil Fertility Management for Lemon Grass Cultivation in Saline Environments

Soil fertility plays a critical role in the growth and productivity of lemongrass (*Cymbopogon citratus*), particularly in saline environments where soil quality is often compromised by high levels of salt. Effective soil fertility management practices are essential to improve the soil's physical, chemical, and biological properties, thus ensuring optimal growth and yield of lemongrass. In the Indian context, where saline soils are prevalent, especially in regions like Gujarat, Rajasthan, and Tamil Nadu, addressing soil fertility challenges is a priority.

Organic Amendments: The addition of organic amendments, such as compost, farmyard manure (FYM), and green manure, has been shown to enhance soil fertility and improve plant growth in saline soils. These amendments help in increasing soil organic matter (SOM), which improves soil structure, water retention, and nutrient availability (Sharma et al., 2006). A study conducted in Rajasthan indicated that the application of FYM at 10 tons per hectare resulted in a 20-25% increase in lemongrass yield under saline conditions (Kumar et al., 2008). Organic matter also aids in reducing the impact of salinity by improving the cation exchange capacity (CEC) of the soil, which is critical for nutrient uptake.

Gypsum Application: Gypsum is widely used in saline and sodic soils to displace sodium ions and improve soil structure. It enhances the infiltration of water, which reduces salt accumulation around plant roots. Studies have shown that the application of gypsum at a rate of 1-2 tons per hectare improves the growth of lemongrass in saline soils, with a significant increase in essential oil yield (Sharma & Gupta, 2007). The application of gypsum helps reduce the pH of the soil, creating a more favorable environment for plant growth.

Table 4: Effect of Soil Amendments on Lemongrass Growth in Saline Soils

Soil Amendment	Lemongrass Biomass Yield (kg/ha)	Essential Oil Yield (L/ha)	Soil pH
No Amendment	2,500	30	8.2
FYM (10 tons/ha)	3,100	35	7.9
Gypsum (1 ton/ha)	3,300	38	7.5
FYM + Gypsum (1 ton/ha)	3,800	42	7.3

Source: Kumar et al., 2008

As shown in Table 4, the combination of organic and inorganic amendments significantly improves lemongrass growth in saline soils. The highest increase in biomass and essential oil yield was observed with the combined application of FYM and gypsum, leading to improved soil pH and fertility.

Micronutrient Fertilization: The application of micronutrients, particularly zinc, iron, and manganese, is essential for lemongrass in saline environments. These micronutrients help in mitigating the adverse effects of salinity stress by enhancing enzymatic activities and improving plant metabolism. Studies have indicated that the foliar application of micronutrient mixtures can result in up to a 15% increase in lemongrass yield in saline areas (Yadav et al., 2007).

In conclusion, managing soil fertility through the use of organic amendments, gypsum, and micronutrient fertilization is crucial for optimizing lemongrass production in saline environments. These practices help improve soil health, reduce salinity stress, and enhance overall productivity.

8. Water Management Techniques for Lemon Grass in Saline Environments

Water management is a critical aspect of lemongrass (*Cymbopogon citratus*) cultivation in saline environments. Salinity stress significantly hampers water availability and uptake by plants, leading to reduced growth and yield. Therefore, adopting effective water management practices is essential to optimize water use and minimize salinity-related stress, especially in arid and semi-arid regions of India where saline soils are prevalent.

Drip Irrigation: Drip irrigation is one of the most efficient methods for water management in saline soils. It delivers water directly to the root zone, minimizing water wastage and ensuring that the plants receive adequate moisture while reducing the spread of salts on the soil surface. According to a study conducted in Rajasthan, the use of drip irrigation in lemongrass cultivation led to a 25-30% increase in yield compared to conventional flood irrigation (Sharma & Kumar, 2007). This method also reduces the risk of salt accumulation around the roots, thereby improving the plant's ability to tolerate salinity stress.

Leaching: Leaching is the process of applying excess water to flush out soluble salts from the root zone. This practice is particularly effective in areas where the salinity level of irrigation water is high. In Gujarat, research has shown that leaching fractions of 15-20% are necessary to maintain acceptable soil salinity levels for lemongrass growth (Sivapalan et al., 2008). Leaching helps in improving soil structure and promoting better nutrient uptake by plants, which in turn enhances plant growth and essential oil production.

Rainwater Harvesting: In regions where rainfall is erratic, rainwater harvesting can be an effective strategy to supplement irrigation. Collecting and storing rainwater allows farmers to use fresh, low-salinity water during dry periods, reducing the dependency on saline groundwater. This approach has been successfully implemented in Tamil Nadu, where rainwater harvesting systems have led to a 20% reduction in irrigation costs for lemongrass farmers (Kumar et al., 2009). By utilizing rainwater, farmers can minimize the salt stress on plants, leading to healthier crops and improved yields.

Water Use Efficiency (WUE): Improving water use efficiency (WUE) is crucial for sustainable agriculture in saline environments. Techniques such as scheduling irrigation based on evapotranspiration rates and using soil moisture sensors can help optimize water use. In a study conducted in Uttar Pradesh, WUE was found to improve by 18% in lemongrass fields when irrigation was scheduled based on real-time soil moisture data (Singh et al., 2006). This approach ensures that water is used efficiently, reducing excess irrigation and preventing salt buildup.

In conclusion, effective water management techniques such as drip irrigation, leaching, rainwater harvesting, and improved WUE are essential for successful lemongrass cultivation in saline environments. These practices not only help in mitigating the effects of salinity but also contribute to sustainable agricultural practices in water-scarce regions of India.

Conclusion

In conclusion, optimizing agronomic practices for enhancing lemongrass (*Cymbopogon citratus*) yield in saline environments requires a multifaceted approach. Soil fertility management, water management techniques, appropriate irrigation systems, and the use of organic and inorganic amendments play vital roles in improving growth and productivity. Organic amendments such as farmyard manure and gypsum, along with micronutrient application, have proven to be effective in mitigating salinity stress and improving soil health. Similarly, efficient water management practices, particularly drip irrigation and leaching, can significantly enhance water use efficiency, minimize salt accumulation, and reduce the impact of saline stress on the crop.

Additionally, the importance of tailored agronomic practices, such as precise fertilizer application and optimal sowing techniques, cannot be overstated. These practices help create favorable growth conditions for lemongrass, particularly in regions where saline soils are widespread, such as Rajasthan, Gujarat, and Tamil Nadu. The integration of these practices has the potential to improve both the biomass and essential oil yields, contributing to better economic returns for farmers.

However, achieving sustainable and high-yielding lemongrass cultivation in saline areas requires continuous research and adaptation of local practices. The adoption of modern technologies, along with traditional knowledge, will be key in addressing the challenges posed by salinity. By focusing on optimizing agronomic practices and employing integrated soil and water management strategies, the potential for lemongrass cultivation in saline environments can be fully realized, ensuring both environmental sustainability and economic profitability for farmers.

This review highlights the necessity for region-specific, scientifically-backed interventions to achieve improved lemongrass yields and ensure the long-term success of this valuable crop in saline-prone areas of India.

References

1. Bajwa, M. S., & Sharma, A. (2006). Enhancing the productivity of lemon grass through improved agronomic practices in saline soils. Indian Journal of Agronomy, 51(1), 52-57.
2. Bhattacharyya, P., Singh, D. P., & Sharma, P. (2006). Effect of soil salinity on the growth of *Cymbopogon citratus* (lemongrass). *Indian Journal of Agronomy, 51*(3), 190-193.
3. Chandran, S., & Radhakrishnan, P. (2009). Organic amendments and their effects on soil fertility and lemongrass yield under saline conditions. Indian Journal of Agricultural Sciences, 79(8), 602-607.
4. Chauhan, B. S., Singh, R., & Sharma, S. (2006). Lemongrass: A promising aromatic plant for essential oil production. *Indian Journal of Essential Oils and Medicinal Plants, 12*(3), 140-145.
5. Chauhan, D. S., & Singh, S. R. (2009). Agronomic practices for optimizing lemongrass yield. *Indian Journal of Horticulture, 66*(4), 437-441.
6. Dhawan, R., & Singh, M. (2007). Effect of irrigation and nutrient management on the growth and yield of lemongrass under saline conditions. Haryana Journal of Agricultural Sciences, 36(4), 238-242.

7. Ghosh, P., & Maiti, S. (2005). Role of soil amendments in the management of salinity in lemongrass cultivation. Indian Journal of Soil and Water Conservation, 3(2), 137-141.
8. Gupta, V. K., & Chauhan, R. (2007). Managing soil salinity for increased productivity in lemongrass farming. Indian Journal of Agronomy, 52(2), 163-167.
9. Kumar, R., Singh, P., & Joshi, R. (2009). Water management practices for sustainable agriculture in saline regions of Tamil Nadu. Indian Journal of Agricultural Sciences, 79(4), 264-268.
10. Munns, R. (2002). Comparative physiology of salt and water stress. *Plant, Cell & Environment, 25*(2), 239-250.
11. Patil, S. D., & Patil, V. D. (2006). Use of gypsum and organic manure in enhancing the yield of lemongrass in saline soil conditions. Indian Journal of Agricultural Sciences, 76(6), 332-335.
12. Rattan, R. K., Singh, S., & Yadav, S. (2008). Irrigation management in saline soils for improved crop yields. *Indian Journal of Soil Science, 57*(2), 142-148.
13. Sharma, A., & Mehta, R. K. (2007). Enhancing crop productivity in saline soils through agronomic practices: A case study of lemongrass cultivation. Journal of Environmental Biology, 28(5), 907-912.
14. Sharma, R., & Kumar, P. (2007). Drip irrigation for effective water management in arid regions. Journal of Irrigation Science, 23(2), 128-133.
15. Singh, A., Sharma, V., & Yadav, S. (2006). Water use efficiency in lemongrass cultivation under saline conditions in Uttar Pradesh. Indian Journal of Agricultural Research, 40(3), 210-214.
16. Singh, M., Singh, R. K., & Chauhan, R. P. (2008). Effect of soil amendments on growth and yield of *Cymbopogon citratus* in saline soils. *Indian Journal of Agronomy, 53*(2), 144-147.
17. Sivapalan, M., Singh, N., & Chauhan, R. (2008). Leaching requirements for managing soil salinity in Gujarat. Indian Journal of Soil Science, 56(1), 73-78.
18. Yadav, J. S., Ram, K., & Choudhary, R. (2007). Salinity stress in Indian agriculture: An overview. *Agricultural Reviews, 28*(2), 138-142.
19. Yadav, R. K., & Bansal, P. K. (2008). Influence of irrigation systems on the growth and yield of lemongrass in saline regions. Agricultural Water Management, 95(2), 170-175.