Journal of Advances in Developmental Research (IJAIDR)
E-ISSN: 0976-4844 ● Website: www.ijaidr.com ● Email: editor@ijaidr.com

The Green Apocalypse: Invasive Plants, Biodiversity Loss and Climate Change

Saju Abraham

Assistant Professor
Department of Botany
Newman College, Thodupuzha, Idukki, Kerala

Abstract

Invasive plant species have emerged as one of the most crusading yet underappreciated drivers of global biodiversity loss, threatening the stability and resilience of ecosystems across biomes. Named as "green apocalypse," the unchecked spread of these species disrupts native plant communities, alters soil chemistry, modifies hydrological cycles, and creates imbalances in food webs. Unlike native flora, invasive plants often possess competitive advantages such as rapid growth, prolific seed production and resistance to local pests and diseases. These traits allow them to outcompete indigenous vegetation, leading to homogenization of landscapes and the collapse of local biodiversity. The consequences are particularly severe in fragile ecosystems, including wetlands, islands and tropical forests, where endemism is high and species have limited adaptive capacity. The impacts of invasive plants extend beyond plant communities, affecting fauna that depend on native species for food and habitat. Pollinators, herbivores and soil organisms are among the most affected, leading to cascading ecological consequences. In addition, the combined effects of climate change and invasive plants will open new avenues of biodiversity loss and species extinctions alarmingly in a warming world. Thus, the paper is an attempt to explore the green apocalypse via the triple action of invasive plants, biodiversity loss and climate change and associated ecological crisis. In short, invasive plant species are among the leading drivers of global biodiversity loss, disrupting ecosystems, outcompeting native flora and altering habitat structures.

Keywords: Green Apocalypse, Biodiversity, Invasive Plants, Climate Change

Introduction

In recent decades, the spread of invasive plant species has emerged as a significant yet often underestimated threat to global biodiversity and ecological balance (CBD, 2009). Invasive species, introduced either intentionally or accidentally beyond their native ranges, are capable of rapidly colonizing new environments, outcompeting native flora and reshaping entire ecosystems with far reaching implications (EEA, 2013). The concept referring to a potential ecological catastrophe where the planet is destroyed or severely damaged due to anthropogenic actions, primarily environmental destruction is referred to as green apocalypse. Invasive plants, with their aggressive growth patterns, high reproductive rates, and adaptability, can alter soil chemistry, disrupt nutrient cycles, change fire regimes and reduce water availability. Such changes compromise the survival of indigenous plant and animal species, leading to homogenization of ecosystems and loss of unique ecological communities.

The problem becomes even more critical when viewed in the context of climate change. Warming temperatures, changing precipitation patterns and more frequent extreme weather events create conditions privileging the spread of invasive plants over native species. As a result, climate change and biological invasions act in synergy, accelerating habitat degradation and reducing the adaptive capacity of ecosystems (Burgiel and Muir, 2010). This interconnected threat not only undermines biodiversity but also weakens ecosystem services such as carbon sequestration, soil stabilization and pollination, which are essential for human survival and for climate regulation itself.

Furthermore, invasive species have direct socio-economic impacts, including reduced agricultural productivity, increased management costs and threats to water resources and human health. Species like *Lantana camara*, *Parthenium hysterophorus*, and *Eichhornia crassipes* have invaded millions of hectares globally, severely affecting both biodiversity and livelihoods (Bhatt *et al.,* 2012). In tropical and subtropical regions such as South Asia and Africa, where biodiversity is rich and communities heavily depend on natural resources, the consequences are especially severe. Despite the magnitude of the issue, invasive plants remain a relatively low priority in environmental policy and conservation planning. Thus, the paper seeks to explore the complex and interlinked relationships between invasive plants, biodiversity loss and climate change. Understanding the role of invasive species in driving ecological disruption is essential not only for conserving biodiversity but also for building resilient ecosystems capable of withstanding the challenges of a rapidly changing climate.

Mechanisms of biodiversity loss caused by invasive plants

Native habitats and the species that depend on them are seriously threatened by invasive plant species, which are acknowledged as one of the main causes of biodiversity loss worldwide. Invasive plants, which have been purposefully or unintentionally brought outside of their natural ranges, frequently have biological traits that provide them with an advantage over native plants. Many invasive species grow faster, form dense canopies or develop extensive root systems that deprive native species of the resources required for their survival, ultimately leading to their local extinction. The key mechanism is the alteration of habitat structure and composition. Invasive plants can fundamentally change the physical characteristics of ecosystems, thereby disrupting the ecological niches that native species have adapted to over evolutionary time. Mooney and Cleland (2001) clearly promulgated the evolutionary impact of invasive species. This disruption leads to the decline of specialized native plants and animals that rely on specific habitat conditions. Moreover, invasive plants can modify soil chemistry by altering nutrient cycling, pH levels, and microbial communities, making it difficult for native species to establish and persist (Weidenhamer and Callaway, 2010). Some invasive species, such as certain legumes, can increase nitrogen levels in soils, benefiting themselves while disadvantaging native plants adapted to nutrient-poor conditions (Wang *et al.,* 2015). Fig 1 depicts various ,mechanisms involved in plant invasions.



Fig 1: Mechanism of plant invasion

In addition to resource competition and habitat alteration, invasive plants also interfere with native species through allelopathy, the release of chemical compounds into the environment that inhibit the germination and growth of neighbouring plants (Orr *et al.,* 2005). This biochemical warfare gives invasive species a powerful advantage in monopolizing space and resources. Furthermore, invasive plants can disrupt mutualistic relationships such as pollination and seed dispersal, which are critical for the reproduction and survival of many native species. By altering the availability of floral resources or changing the behaviour of pollinators, invasive plants reduce the reproductive success of native plants, further driving biodiversity decline. Fire regime modification is yet another pathway through which invasive plants contribute to biodiversity loss (Brooks *et al.,* 2004). Certain invasive grasses and shrubs increase the frequency, intensity or spread of wildfires, creating conditions that favour fire-adapted invasive species while simultaneously destroying native plant communities that may not be resilient to frequent fires. This feedback loop of invasion and disturbance leads to the progressive homogenization of ecosystems, eroding both species richness and ecological complexity. Collectively, these mechanisms demonstrate that the threat posed by invasive plants is multifaceted, operating at genetic, species and ecosystem levels.

Trophic cascades and faunal decline

Trophic cascades represent one of the most profound ecological processes through which disruptions in food webs lead to widespread changes in ecosystem structure, function and biodiversity. These cascades typically occur when alterations at the top of the food chain, such as the removal or decline of apex predators trigger a chain of indirect effects that move down through the trophic levels, ultimately impacting vegetation, nutrient cycles and entire habitats. The decline of faunal populations, especially key herbivores carnivores, and omnivores, plays a central role in initiating or accelerating trophic cascades. Walsh *et al.* (2016) reported that invasive species triggers a massive loss of ecosystem services through trophic cascade which are observed across terrestrial, freshwater and marine ecosystems, highlighting the interconnectedness of ecological relationships and the fragility of biodiversity when keystone species are lost.

One of the primary ways trophic cascades lead to faunal decline is through the disruption of predator-prey relationships *via* invasive alien plants. According to Doherty *et al.* (2015) the interactions between invasive predators and other ecological disturbances along with invasive alien plants will have catastrophic ecological ramifications. The loss of key herbivorous species by the influence of invasive plants also initiates trophic cascades by disrupting plant-animal interactions such as seed dispersal and pollination. Large frugivores like elephants, primates and birds play vital roles in dispersing seeds across landscapes, facilitating forest regeneration and maintaining plant diversity. Declines in these species often result in changes in forest composition, favouring plant species with alternative dispersal mechanisms or those that do not depend on large animals for reproduction. These shifts reduce habitat heterogeneity, ultimately affecting the broader faunal community by limiting the availability of diverse food resources and nesting sites. These cascading effects underscore the critical need to conserve entire food webs rather than focusing solely on individual species. Understanding and managing trophic interactions is essential for maintaining biodiversity, ensuring ecological stability and preserving the ecosystem services upon which human societies rely.

Socioeconomic implications

Invasive alien plant species pose not only ecological threats but also significant socioeconomic challenges that affect livelihoods, public health, agricultural productivity and the overall well-being of communities (Senator and Rozenberg, 2007). These species, once introduced outside their native habitats, establish and spread rapidly, disrupting natural ecosystems and altering land use patterns. The socioeconomic implications of such invasions are profound, particularly in regions where local economies and subsistence livelihoods are closely tied to natural resources, agriculture and ecosystem services. One of the primary economic impacts of invasive plants is the reduction in agricultural productivity (Reaser *et al.,* 2007). Many invasive species aggressively colonize farmlands, grazing areas and irrigation channels, competing with crops for nutrients, water and sunlight. Plants such as *Parthenium hysterophorus* and *Chromolaena odorata* have become notorious for invading cultivated fields and pastures in several parts of the world, including South Asia and Africa (Koutika and Rainey, 2010; Kumari *et al.,* 2014). Their rapid spread often leads to decreased crop yields, forcing farmers to spend more on weeding, herbicides and land restoration efforts which all adds financial burden particularly on smallholder farmers. Additionally, invasive plants reduce the carrying capacity of grazing lands by replacing palatable forage species, thereby impacting livestock production and rural livelihoods dependent on animal husbandry.

Water resources are also significantly affected by invasive plants. Species like *Eichhornia crassipes* (water hyacinth) form dense mats on the surface of lakes, rivers and reservoirs, obstructing water flow, hampering fishing activities and increasing water evaporation rates (Gichuki *et al.,* 2012). These infestations can clog irrigation systems, reduce hydroelectric power generation and lead to increased costs in maintaining water infrastructure. In regions where water availability is already a concern, such invasions pose direct threats to agricultural sustainability, drinking water supplies and energy production (Rocha-Ramírez *et al.,* 2014; Tewabe, 2015). The socioeconomic implications extend to public health as well. Some invasive plants release allergens or toxic substances, causing respiratory problems, skin irritations or even facilitating the breeding of disease vectors like mosquitoes in stagnant water bodies infested with aquatic invasive plants. The health burden created by such invasions strains already limited healthcare resources, especially in rural or developing regions.

Invasive plants and public health

Invasive plant species are often discussed in the context of environmental degradation and biodiversity loss, but their impacts on public health are equally significant and frequently overlooked. These impacts can manifest in various direct and indirect ways, affecting respiratory health, skin conditions, water quality, food security and mental well-being (Kopelson, 2014). As invasive plants spread across ecosystems, they often alter the living conditions of both humans and other organisms, creating new challenges for public health systems, particularly among vulnerable rural and peri-urban communities. One of the most direct health impacts of invasive plants is through allergenic reactions. Several invasive species, such as *Parthenium hysterophorus*, release pollen and airborne allergens that can trigger respiratory diseases, including allergic rhinitis, asthma and bronchitis (Gajanan and Fernandes, 2015). Prolonged exposure to these allergens, especially during flowering seasons, can significantly reduce the quality of life for affected individuals. Skin contact with certain invasive species can also lead to dermatitis and other allergic skin reactions, creating chronic irritation and discomfort, especially among populations that rely on natural resources for their livelihoods (Otang *et al.,* 2014).

Beyond respiratory and skin-related ailments, invasive aquatic plants such as *Eichhornia crassipes* pose severe risks to water quality and sanitation. Dense mats of water hyacinth provide ideal breeding grounds for mosquitoes which directly contributes to the spread of mosquito-borne diseases particularly in tropical and subtropical regions. The proliferation of invasive plants in aquatic ecosystems also reduces access to clean water for drinking, bathing, and irrigation, compounding existing health risks associated with waterborne diseases. Mental health impacts associated with invasive plant infestations are another emerging concern. The degradation of landscapes, loss of traditional livelihoods, and increased burdens on daily life caused by invasive plants can lead to stress, anxiety and depression, particularly in rural communities where identity and cultural practices are closely tied to land and natural resources. The psychological burden of managing relentless plant invasions, combined with economic losses, can further strain already vulnerable populations.

Conclusion

The rapid spread of invasive plant species across the globe has ushered in what can only be described as a "Green Apocalypse", a silent ecological crisis where aggressive non-native flora systematically dismantle native ecosystems, accelerate biodiversity loss and exacerbate the impacts of climate change. As the paper has demonstrated, invasive plants do not merely occupy space; they actively reshape ecosystems through competitive exclusion, habitat modification, trophic disruptions and allelopathic effects often with irreversible consequences. The compounding effects of climate change further intensify this crisis by creating favourable conditions for invasives to thrive while weakening the resilience of native species. The result is a dangerous feedback loop: as invasive plants dominate landscapes, they reduce biodiversity, which in turn diminishes capacity of ecosystem to adapt to climatic stressors, leaving ecosystems more vulnerable to further invasions. The ecological ramifications of this phenomenon are profound and multifaceted. Moreover, invasive plants have demonstrated their ability to alter soil chemistry, disrupt hydrological cycles and even modify fire regimes. These changes do not occur in isolation; they cascade through food webs, threatening pollinators, herbivores and predators alike. The decline of specialist species, those reliant on specific native plants is particularly alarming, as their loss signals the unravelling of intricate ecological networks that have evolved over millennia. Beyond ecology, the socioeconomic costs of invasive plants are staggering. The Green Apocalypse is not an inevitable fate but an imminent crisis, fuelled by globalization, land-use changes and a historical underestimation of invasive species' impacts. The stakes could not be higher: biodiversity loss and climate change are existential threats and invasive plants sit at the intersection of both. The time to act is now; the alternative is a future where biodiversity is irretrievably eroded and the ecological services the world depend on are forever diminished.

References

1. Bhatt, J.R., Singh, J.S., Singh, S.P., Tripathi, R.S. and Kohli, R.K. 2012. *Invasive alien plants: an ecological appraisal for the Indian subcontinent*. CABI Digital Library. ISBN 978-1-84593-907-6.
2. Brooks, M.L., D'Antonio, C.M., Richardson, D.M., DiTomaso, J.M., Grace, J.B., Hobbs, R.J., Keeley, J.E., Pellant, M. and Pyke, D. 2004. Effects of invasive alien plants on fire regimes. *BioScience*. 54(7): 677-688.
3. Burgiel, S.W. and Muir, A.A. 2010. *Invasive Species, Climate Change and Ecosystem-Based Adaptation: Addressing Multiple Drivers of Global Change*. Global Invasive Species Programme September 2010. doi:10.13140/2.1.1460.8161.
4. CBD. 2009. *Invasive Alien Species a threat to biodiversity*. Secretariat of the Convention on Biological Diversity, Quebec, Canada. ISBN: 92-9225-119-8.
5. Doherty, T.S., Dickman, C.R., Nimmo, D.G. and Ritchie, E.G. 2015. Multiple threats, or multiplying the threats? Interactions between invasive predators and other ecological disturbances. *Biological Conservation*. 190:60-68.
6. EEA. 2013. *Invasive alien species: a growing problem for environment and health*. https://www.eea.europa.eu/highlights/invasive-alien-species-a-growing.
7. Gajanan, G. and Fernandes, N. 2015. Parthenium Allergy in Indian Population: Clinical Significance. *Journal of Evidence based Medicine and Healthcare*. 2(41):7109-7117.
8. Gichuki, J., Omondi, R., Boera, P., Okorut, T., Matano, A.S., Jembe, T. and Ofulla, A. 2012. Water hyacinth *Eichhornia crassipes* (Mart.) Solms-Laubach dynamics and succession in the Nyanza Gulf of Lake Victoria (east Africa): implications for water quality and biodiversity conservation. *Scientific World Journal*. 2012:106429. doi: 10.1100/2012/106429.
9. Kopelson, H. 2014. The Impacts of Invasive Plant Species on Human Health. City Wild. Paper 6. http://digitalcommons.esf.edu/citywild/6.
10. Koutika, L.S. and Rainey, H.J. 2010. *Chromolaena odorata* in different ecosystems: Weed or fallow plant? *Applied Ecology and Environmental Research*. 8(2):131-142.
11. Kumari, P., Sahu, P., Soni, M. and Awasthi, P. 2014. Impact of *Parthenium hysterophorus* L. Invasion on Species Diversity of Cultivated Fields of Bilaspur (C.G.) India. *Agricultural Sciences*. 5:754-764.
12. Mooney, H.A. and Cleland, E.E. 2001. The evolutionary impact of invasive species. *Proc. Natl. Acad. Sci. U.S.A.* 98(10):5446-5451.

13. Orr, S., Rudgers, J. and Clay, K. 2005. Invasive Plants can Inhibit Native Tree Seedlings: Testing Potential Allelopathic Mechanisms. *Plant Ecol.* 181:153-165.
14. Otang, W.M., Grierson, D.S. and Afolayan, A.J. 2014. A survey of plants responsible for causing irritant contact dermatitis in the Amathole district, eastern cape, South Africa. *Journal of Ethnopharmacology*. 157:274-284.
15. Reaser, J., Meyerson, L., Cronk, Q., De Poorter, M., Eldrege, L., Green, E., Kairo, M., Latasi, P., Mack, R.N., Mauremootoo, J., O'Dowd, D., Orapa, W., Sastroutomo, S., Saunders, A., Shine, C., Thrainsson, S. and Vaiut, L. 2007. Ecological and socioeconomic impacts of invasive alien species in island ecosystems. *Environmental Conservation*. 34(2):98-111.
16. Rocha-Ramírez, A., Robles-Valderrama, E. and Ramírez-Flores, E. 2014. Invasive alien species water hyacinth *Eichhornia crassipes* as abode for macroinvertebrates in hypertrophic Ramsar Site, Lake Xochimilco, Mexico. *Journal of Environmental Biology*. 35:1071-1080.
17. Senator, S.A. and Rozenberg, A.G. 2017. Assessment of economic and environmental impact of invasive plant species. *Biol. Bull. Rev.* 7:273-278.
18. Tewabe, D. 2015. Preliminary Survey of Water Hyacinth in Lake Tana, Ethiopia Tana, Ethiopia. *Glob. J. Allergy*. 1(1):013-018.
19. Walsh, J.R., Carpenter, S.R. and Vander Zanden, M.J. 2016. Invasive species triggers a massive loss of ecosystem services through a trophic cascade, *Proc. Natl. Acad. Sci. U.S.A.* 113(15):4081-4085.
20. Wang, C., Xiao, H., Liu, J., Wang, L. and Du, D. 2015. Insights into Ecological Effects of Invasive Plants on Soil Nitrogen Cycles. *American Journal of Plant Sciences*. 6:34-46.
21. Weidenhamer, J.D. and Callaway, R.M. 2010. Direct and indirect effects of invasive plants on soil chemistry and ecosystem function. *J. Chem. Ecol*. 36(1):59-69.